



Murdoch
UNIVERSITY

MURDOCH RESEARCH REPOSITORY

This is the author's final version of the work, as accepted for publication following peer review but without the publisher's layout or pagination. The definitive version is available at

<http://dx.doi.org/10.1071/AM11040>

Wilson, B.A., Valentine, L.E., Reaveley, A., Isaac, J. and Wolfe, K.M. (2012) Terrestrial mammals of the Gnangara Groundwater System, Western Australia: history, status, and the possible impacts of a drying climate. Australian Mammalogy, 34 (2). pp. 202-216.

<http://researchrepository.murdoch.edu.au/10909/>

Copyright © Australian Mammal Society 2012

It is posted here for your personal use. No further distribution is permitted.

**Terrestrial mammals of the Gnamptara Groundwater System, Western Australia:
history, status, and the possible impacts of a drying climate.**

Barbara A. Wilson^a, Leonie E. Valentine^{a,b}, Alice Reaveley^a, Joanne Isaac^a and
Kristen M. Wolfe^a

^a Gnamptara Sustainability Strategy, Department of Environment and Conservation,
PO Box 1167, Bentley Delivery Centre, WA 6983.

^b WA State Centre of Excellence for Climate Change, Woodland and Forest Health,
Murdoch University, South St, Murdoch WA 6150.

Corresponding author email address: Barbara.Wilson@dec.wa.gov.au

Abstract

Over the last 30 years declining rainfall and increased aquifer abstraction have heavily impacted water availability and ecosystems on the Gnamptara Groundwater System (GGS). The mammal fauna of the area is considered to have been rich, with up to 28 terrestrial and 5 volant native species recorded since European settlement. This study investigated previous and current distribution of mammals on the GGS, and assessed potential impacts of predicted rainfall and groundwater declines on mammals. A general survey was conducted at 40 sites, and targeted trapping was undertaken for *Hydromys chrysogaster* and *Isoodon obesulus fusciventer* at wetlands. Nine native and seven introduced terrestrial mammal species were recorded during the general survey and capture rates were very low (1.05%). The most commonly captured native species was *Tarsipes rostratus*. There is evidence that only 11 (9 recorded and 2 considered to be extant) of the 28 historically recorded terrestrial native mammals are still persisting in the area. The species predicted to be most susceptible to rainfall and groundwater level declines include *H. chrysogaster*, *I. obesulus fusciventer*, and *T. rostratus*. Management and recovery actions required to protect mammals under predicted climatic changes include identification and maintenance of refugia and ecological linkages, supplementation of lakes, development of ecologically appropriate fire regimes, and control of predators.

Additional keywords:

Western Australia, Northern Swan Coastal Plain, aquifer, ecological linkages, fire regimes, groundwater dependency.

Introduction

In the past decade numerous studies have demonstrated the impacts of climatic changes in a variety of ecosystems and communities around the world (Hughes, 2000; Walther *et al.*, 2002; Parmesan and Yohe, 2003). In Mediterranean systems, such as south-western Australia, climate change predictions include lower rainfall and extended drought periods (IPCC, 2007). If correct, these changes to the climate are likely to pose significant threats to vulnerable mammal species, especially in fragmented ecosystems.

Low rainfall has been shown to have a significant influence on the abundance of small mammals, including brush-tailed phascogales (*Phascogale tapoatafa*) (Rhind and Bradley, 2002) and honey possums (*T. rostratus*) (Wooller *et al.*, 1998; Bryant, 2004; Bradshaw *et al.*, 2007) in south-western Australia. There is also recent evidence that low rainfall and drought have severe impacts on the abundance of species such as agile antechinus (*Antechinus agilis*) and swamp antechinus (*A. minimus*) in south-eastern Australia (Wilson *et al.*, 2007; Parrott *et al.*, 2007; Sale *et al.*, 2008; Magnúsdóttir *et al.*, 2008). Although there has been little work on the relationship of mammals and groundwater levels, Braithwaite and Muller (1997) found strong correlations between mammal population declines and decreased groundwater levels in an area of the Australian wet-dry tropics. In contrast, Woinarski *et al.* (2001; 2010) did not find a direct link between rainfall and mammal abundance at the same site, and suggest that declines in mammal abundance may be due to other threatening processes such as a high frequency of fire.

The record of extinction of mammals in Australia since European settlement is recognised as being higher than elsewhere in the world (Burbidge and McKenzie, 1989; Morton, 1990; Short and Smith, 1994; Maxwell *et al.*, 1996). A number of disturbance factors have been identified as contributors to the extinctions and declines including land clearance and habitat fragmentation, introduced predators, changed fire regimes, disease and rainfall decline (Burbidge and McKenzie, 1989; Morton, 1990; Maxwell *et al.*, 1996; Johnson, 2006). McKenzie *et al.* (2007) found that the loss of

mammal species was greater in arid regions compared to high rainfall regions. Most of the region-to-region variation in attrition was attributed to variables such as mean annual rainfall, environmental change, phylogenetic similarity, and body-weight distribution. Mammals that occur in areas predicted to have reduced rainfall and ground water levels are thus highly susceptible to population declines.

The IPCC (2007) has identified a number of regions (coastal) and ecosystems (wetland and Mediterranean) where faunal biodiversity is likely to be at particularly high risk from the impacts of climate change. South-western Australia is predicted to undergo the most intense drying of any region in Australia by 2100 (Pitman and Perkins, 2008), with modelling suggesting that rainfall will decline by as much as 20% by 2030 and 60% by 2070 (Jones and Preston, 2006). The declines in rainfall will predominantly occur during winter months, although declines in spring and summer rainfall are also expected. In addition to direct impacts of declining rainfall on mammals, there are threats from indirect impacts of declining aquifers. The ecological impacts of groundwater declines on groundwater dependent ecosystems (GDEs), including surface terrestrial and aquatic systems, have been recognised as significant (Eamus *et al.* 2006; Nevill *et al.* 2010).

The Gnamptara Groundwater System (GGS), which is located in south-western Australia on the Swan Coastal Plain (SCP), is an area at particularly high risk to rainfall declines (Malcolm *et al.*, 2006). The system covers approximately 220 000 ha and provides the city of Perth with approximately 60% of its drinking water (Figure 1). The groundwater system consists of an unconfined, superficial aquifer known as the Gnamptara Mound (Government of Western Australia, 2009). In the past 30 years, a drying climate has been a significant feature of the GGS. Rainfall has declined largely compared to the long-term average (Figure 2) and, together with increased aquifer abstraction, have strongly influenced the groundwater storage (Figure 3), resulting in groundwater levels that have decreased by up to 4 m in some areas (Yesertener, 2007; Vogwill *et al.*, 2008). It is predicted that in the future there could be an even greater decline in rainfall (IOCI, 2005; Jones and Preston, 2006; Preston and Jones, 2006) and under all but the most optimistic assumptions for climate, groundwater levels will decline (Vogwill *et al.*, 2008).

Declining groundwater levels have emerged as a significant threat to wetland-associated vertebrate fauna across the SCP and in the GGS area. The wetlands are some of the most biologically productive areas of the SCP and directly or indirectly support most of the wildlife (Horwitz *et al.* 2009). Declines in wetlands are likely to have significant impacts on mammal fauna and their habitats; however, there is little knowledge of both the habitat components that are most affected, and the communities or taxa that are most susceptible to declining groundwater levels. Further, there is strong evidence of a progressive reduction in canopy foliage cover and change in floristic composition of the *Banksia* woodlands on the GGS as the regional water table has declined (Sommer and Froend 2011).

The mammal fauna of the northern SCP that covers the GGS area is considered to have been historically rich, with up to 33 native species recorded since European settlement (Kitchener *et al.*, 1978). However, in the 1970s only 12 species were recorded across large contiguous and relatively undisturbed vegetation in the north of the GGS area (Kitchener *et al.*, 1978). Subsequent studies in inner urban fragments (1 - 338 ha in size) found few mammals surviving with only seven native and six introduced species recorded (How and Dell, 1993; 1994; 2000). However, a comprehensive fauna survey has not been conducted in the large remnants since the late 1970s and the current status and persistence of mammals is unclear (Mitchell *et al.* 2003).

In this paper we assess the historic and current occurrence and distribution of terrestrial mammal fauna across the GGS, and assess the susceptibility of taxa to declining rainfall and groundwater levels. We consider the implications for conservation, recovery and management under a drying climate.

Materials and methods

Study area

The area covered by the GGS is located on the SCP and represents a distinct water catchment that extends from Perth (Swan River) in the south, to the Moore River and Gingin Brook in the north, and from the Darling Scarp in the east to the Indian Ocean in the west (Figure 1) (DOW, 2008; Government of Western Australia, 2009).

Although there have been large amounts of clearing for urbanisation and agriculture,

the total remnant native woodland in the GGS covers more than 100 000 ha and includes the largest continuous area of remnant vegetation on the SCP, south of the Moore River. The distribution of vegetation is predominantly determined by the underlying landforms, soils, depth to water table and climatic conditions (Heddl *et al.*, 1980). The Swan Coastal Plain can be divided into a sequence of broad geomorphic units lying parallel to the coast that are characterised by a progression of aeolian sands in the west to alluvial and/or colluvial deposits in the east (Playford *et al.* 1976). Within the GGS study area, the major landforms include three main dune systems Quindalup, Spearwood and Bassendean dunes (Government of Western Australia 2000; Playford *et al.* 1976). These landform units are composed of a variety of soils and varying surface geology. The younger Quindalup Dunes are close to the coast, followed by the Spearwood Dunes associated with Tamala Limestone ridges within about ten kilometres of the coast, and the inland Bassendean Dunes which are older and flatter, and contain leached and slightly acidic sands. The vegetation is typically a *Banksia* overstorey, with sporadic stands of *Eucalyptus* and *Allocasuarina*, and an understorey consisting mainly of low shrubs from the Myrtaceae, Fabaceae and Proteaceae families. Interspersed among these woodlands are many seasonal damplands, swamps and permanent wetlands, fringed by *Banksia littoralis* and *Melaleuca* trees (Semeniuk *et al.*, 1990).

The GGS experiences a dry Mediterranean-type climate (Beard, 1984), with hot dry summers (December – March) and cool wet winters (June – August), and an annual average of 807 mm. Rainfall and runoff declines in the last 30 years have been substantial, with approximately 21% less rainfall and 64% less runoff between 1997 - 2003 compared to 1911 – 1974 (Yesertener, 2007).

Assessment of the historic occurrences of mammals

Information on previous records and studies was collated from publications, survey team reports and fauna surveys conducted on the GGS, including peer-reviewed journals, state and local government surveys, and university theses (see references listed in Table 1). Records of the historic occurrence and abundance of mammal species were collated and compared to the records for the current study.

Fauna surveys

Fauna surveys were conducted between 2007 and 2008. A general trapping program was conducted on terrestrial mammal species across the study area, as well as two targeted trapping programs for southern brown bandicoot (*I. obesulus fusciventer*) and water rat (*H. chrysogaster*).

General survey trapping

The sampling regime consisted of 40 sites in the major areas of continuous remnant bush land in the northern and eastern areas of the GGS. A stratified sampling procedure was employed to select sites to represent the major landform units (Quindalup, Spearwood and Bassendean), vegetation communities (Banksia woodland, coastal scrub, jarrah forest, tuart forest and Melaleuca wet or damp land). Some categories, such as the Quindalup landform were not well represented due to their limited occurrence and/or poor accessibility in the study area.

The trapping design included an array of pitfall traps (20 L buckets), and a transect line of small aluminium box traps (Elliotts: 9 x 10 x 33 cm) and cage traps (Sheffields: 20 x 20 x 56 cm). Each site contained one pitfall trap array, consisting of 10 pitfall traps arranged in a Y shape with three pitfall traps along each radiating arm and a central pit. Pitfall traps were placed at approximately 7 metre intervals and were connected with 30 cm high aluminium fly wire drift fence. Twenty Elliott and ten cage traps were positioned along a 300 m-long transect between two adjacent sites of pitfall trap arrays. Elliott traps were located at 15 m intervals, and cage traps at 30 m intervals. Both trap types were baited with universal bait comprising a mixture of peanut paste, rolled oats and sardines. Traps were opened for 12 – 20 nights in total for the three trapping periods (spring 2007, autumn 2008 and spring 2008). While open, all traps were checked once per day in the early morning. Captured animals were identified, processed and released onsite. Mammals were ear notched for recapture purposes (*I. obesulus* received an individually numbered ear tag).

Taxonomic nomenclature, including common names, followed the Western Australian Museum protocols. Incidental observations of species were also recorded.

Targeted trapping surveys

A targeted trapping survey for *I. obesulus* was undertaken at nine sites in May 2008 (Figure 1). Sites were located in areas of suitable habitat or based on historical

records in areas that contained a permanent wetland or were swampy or damp for part of the year (T. Friend, M. Bamford, *pers. comm.*, 2007). The targeted trapping survey for *H. chrysogaster* was conducted in May 2008 at permanent lakes considered to provide suitable habitat for the species (Lake Joondalup, Lake Goollellal and Lake Loch McNess).

Transects of cage traps set at 50 m intervals were established in vegetation along the wet or dampland for the target species. Between 10 and 30 traps were set depending on available habitat. Results were standardised to captures per 100 trap nights. Universal bait (a mixture of peanut paste, rolled oats) was used in traps set to capture *I. obesulus* and sardines were used as bait for *H. chrysogaster*. All sites were trapped over 4 nights, except for Lake Goollellal, which was trapped for 3 nights. Captured animals were identified, measured and released onsite. Captured *I. obesulus* individuals were ear-tagged in each ear with unique identification tags. *Hydromys chrysogaster* individuals were ear-notched in one ear.

Assessment of susceptibility of mammals to groundwater and rainfall declines

Information on those mammal species likely to be impacted by decreased aquifer levels and declining rainfall was assessed based on the life history of species and information collated from the literature.

The extent of reliance on groundwater, either in wetland or phreatophytic habitats, was used to classify species into one of three categories (modified from Bamford and Bamford (2003) as:

High: Species that rely on aquatic habitats in wetlands and likely to become locally extinct if surface water disappears.

Strong: Species reliant on wetland vegetation and likely to be impacted depending on changes to riparian vegetation.

Low: Species likely to be indirectly affected.

Further we assessed the potential effect of declining rainfall on the abundance of *T. rostratus* on the GGS using capture rate as a surrogate for population density (see Wooller *et al.*, 1981). First, a linear equation for the relationship between capture rate and the annual rainfall from the previous year was calculated using Wooller *et al.*

(1998) data on capture rates of *T. rostratus* and rainfall from the Fitzgerald River National Park area. These data represent a long-term data set (conducted over 13 years) and establishes a relationship between rainfall and capture rates of *T. rostratus* that is useful for predicting potential change in capture rates in the GGS. We used this linear equation to predict density of *T. rostratus* using rainfall data from the GGS area. We modelled the effect of declining annual rainfall on the predicted density of *T. rostratus* using a number of scenarios, ranging from no decline in annual rainfall to a maximum of 60% decline. Initial values (i.e. assuming no decline) were calculated based on the long-term estimated annual rainfall for the GGS area of 807 mm average from 1905–2007 (Bureau of Meteorology). We then overlaid annual rainfall from recent years (2006 to 2010).

Results

Historic occurrences of mammals in study area

Early collections of mammals from the northern SCP were by John Gilbert in two expeditions (1839-1840 and 1842 -1843), and Shortridge from 1904 to 1907 (Kitchener *et al.*, 1978). These and some other early records resulted in a total of 33 native mammal species, including 5 bat species, as occurring on the northern SCP (Table 1). The Western Australian Museum undertook a comprehensive survey of the extant mammals on the northern SCP in 1977-78 (Kitchener *et al.*, 1978). This survey comprised 24 443 trap nights and confirmed the persistence of only 12 native species (9 ground-dwelling and 3 bat species) of the original 33 species recorded. The areas surveyed included representative communities of the major landforms on the plain including Quindalup, Spearwood, Bassendean and Pinjarra plain soils and were considered to be most likely to still contain extant species as they consisted of large tracts of relatively undisturbed vegetation (e.g. Yeal Nature Reserve, Yanchep National Park) (Kitchener *et al.*, 1978).

Of the original mammal species listed as occurring on the GGS during the last century, 11 species are now gazetted under the WA *Wildlife Conservation Act 1950* as Schedule 1 and one species, the crescent nailtail wallaby (*Onychogalea lunata*), is listed as Schedule 2 (see Table 1 for category definitions). Under the DEC Priority Fauna List (i.e. taxa that have not yet been adequately surveyed to be listed under Schedule 1 or 2), two species are listed as Priority 4 and one as Priority 5 (Table 1).

Eight species are also listed as vulnerable and one species as endangered under the *EPBC Act 1999* (Table 1).

There is little information on the historical abundance and distribution of the mammalian taxa on the northern SCP, although some anecdotal evidence has been compiled by Abbott (2008). Since the late 1970s, a number of species have not been recorded (Table 1).

Western grey kangaroos (*Macropus fuliginosus*) were historically recorded as abundant on the northern SCP and continue to be recorded commonly. They utilise woodland, shrubland and heath formations (Kitchener *et al.*, 1978). Historically western brush wallabies (*Macropus irma*) were frequently sighted from coastal sites in the west to Gingin in the east but appears to have declined since the 1970s (Kitchener *et al.*, 1978). The only population of this species that now occurs on the GGS is in Whiteman Park where fox baiting is undertaken.

Woylies (*Bettongia penicillata ogilbyi*) occurred in the northern SCP (and further north) before disappearing from the area (Maxwell *et al.*, 1996). Historically *I. obesulus* was considered plentiful near Perth; however, by the 1970s it was sparsely distributed on the northern SCP and only two specimens were captured in 1977-88 (Kitchener *et al.*, 1978).

Historically chuditch (*Dasyurus geoffroii*) were common close to the Swan River and specimens in the Western Australian Museum from the northern SCP were collected mainly during the 1900s. This species was not trapped during the 1977-78 Western Australia Museum survey (Kitchener *et al.*, 1978), nor have any been trapped in subsequent fauna surveys. However, there have been intermittent sightings and road kills of *D. geoffroii*, which indicate that the species may persist at a low level in the area (B. Johnson *pers. comm.*, M. Bamford, *pers. comm.*). These individuals may be itinerant individuals moving into the area from populations in the Darling Scarp.

Common brushtail possums (*Trichosurus vulpecula*) were recorded as plentiful on the northern SCP however only one specimen was recorded (Yanchep National Park) in the 1970s survey (Kitchener *et al.*, 1978). The species is likely to still survive in very

low numbers in isolated pockets of vegetation (e.g. Bold Park, Yanchep National Park, Ellenbrook Nature Reserve, Lake Goollellal) and in some north-western suburbs of Perth (How *et al.* 1996, J. Wheeler *pers. comm.*, M. Bamford *pers. comm.*). Kitchener *et al.* (1978) considered that western pygmy possums (*Cercartetus concinnus*) were uncommon across the northern SCP during the 1900s as only 20 specimens were lodged at the museum. The species was not caught in the 1970s survey and since then the only record of their persistence on the GGS is from the Lexia Wetlands area in 2006 (R. Davis, *pers. comm.*). Historically *T. rostratus* was considered uncommon on the northern SCP and only one specimen was recorded in the 1970s survey (Kitchener *et al.*, 1978). However, *T. rostratus* has been recorded infrequently in subsequent fauna surveys in the GGS area since this time (Ecologia Environmental Consultants, 1997). Echidnas (*Tachyglossus aculeatus*) were not collected in the 1977-78 survey and are considered to be uncommon in the south west and on the northern SCP (Kitchener *et al.*, 1978; Abbott, 2008).

The Western Australian sub-species *R. fuscipes fuscipes* is considered to be sparsely distributed on the northern SCP with specimen records limited to the coastal Yanchep area in 1975, and 1977-78 (Kitchener *et al.*, 1978). There are few records of *H. chrysogaster* between Moore River and Swan River on the northern SCP; however this species has been recorded at locations in the GGS area at Lake Jandabup and Lake Joondalup in 1977-78 (Kitchener *et al.*, 1978).

A total of nine introduced species have been recorded in the GGS study area since settlement (dingo (*Canis lupus dingo*), red fox (*Vulpes vulpes*), ferret (*Mustela putorius*), feral cat (*Felis catus*), black rat (*Rattus rattus*), house mouse (*Mus musculus*), European rabbit (*Oryctolagus cuniculus*), feral pig (*Sus scrofa*), and goat (*Capra hircus*).

Current occurrence and distribution of fauna

General survey trapping

The trapping effort (2007-08) included 5 600 pitfall trap nights, 5 600 Elliott trapping nights and 2 800 cage trapping nights. Six mammal species were trapped, four native species (*T. aculeatus*, *Sminthopsis* sp., *T. rostratus*, *R. fuscipes*) and two introduced species (*M. musculus* and *R. rattus*) (Table 2). Mammal capture rates were very low

(1.05%). Captures were higher in pitfalls (1.88%) than cage or Elliott traps (0.50%), although different species were captured in the different trap types. The introduced *M. musculus* was captured at 45% of sites and the native *T. rostratus* at 30% of sites. One *R. fuscipes*, three *T. aculeatus* and one *Sminthopsis* sp. were also captured. The most consistently captured native mammal species was *T. rostratus*. A total of 26 individuals were captured at 14 sites that were widely distributed across the study area (Figure 1). They were captured on Bassendean and Spearwood soils in *Banksia*, Jarrah and *Melaleuca* vegetation types. A further three native species (*T. vulpecula*, *M. fuliginosus*, *M. irma*) and five introduced species were observed (*V. vulpes*, *F. catus*, *O. cuniculus*, *S. scrofa*, *C. hicrus*).

Targeted trapping surveys

The trapping effort for the targeted surveys involved 930 cage trap nights. *Isodon obesulus* was captured at five of the nine sites where trapping was conducted (Table 3). At two sites, Little Badgerup Swamp and Nowergup Nature Reserve, only one individual was captured while the highest number was recorded at Twin Swamps Nature Reserve (Table 3). Non-target native captures included *R. fuscipes* and *T. aculeatus*. *R. fuscipes* was only captured at two sites, but were recorded in greater numbers at Loch McNess. The introduced *R. rattus* and *M. musculus* were captured at most sites and seem to be ubiquitous throughout the area.

Hydromys chrysogaster was captured at all three of the selected sites (Table 3). The highest number of individuals was captured at Lake Goollellal (3 males, 3 females) and Loch McNess (3 males, 3 females), while only one individual was captured at Lake Joondalup (1 female). Non target captures included *R. fuscipes*, *R. rattus* and *M. musculus* (Table 3).

Susceptibility of mammals to declining groundwater levels

Of the terrestrial mammals considered to be extant on the GGS (Table 1), four are highly or strongly susceptible to altered groundwater levels (Table 4).

Hydromys chrysogaster is the only strictly groundwater-dependent mammal and thus is significantly threatened by declining rainfall and associated declines in groundwater level (Morris, 2000; Bamford and Bamford, 2003; Atkinson *et al.*, 2008). *Hydromys*

chrysogaster lives in permanent bodies of fresh or brackish water including lakes, rivers and streams, and the presence of the species in these systems is related to the size of the wetland (Olsen, 1985; Morris, 2000; Smart, 2009). Habitat requirements identified for the species include areas suitable for location of burrows and nests such as steep banks, the presence of logs, rocks or reeds as feeding areas, and adequate riparian and stream vegetation cover (Woollard *et al.*, 1978).

Across the GGS, the occurrence of *I. obesulus* appears to be strongly linked to wetland-associated vegetation (Bamford and Bamford, 1994). In our study, the species were captured at five of the nine targeted survey sites that were located in areas that contained a permanent wetland or were swampy or damp for part of the year. The highest numbers of *I. obesulus* were recorded at Twin Swamps Nature Reserve, which is both baited and fenced to deter foxes.

Historically *R. fuscipes* has only been recorded sporadically across the GGS and were captured at only three sites in our study, which were located either on coastal dune swales or associated with lakes and wetlands. The highest density ($n = 4$) was recorded in wetland vegetation at Loch McNess.

On the GGS *M. irma* prefers *Eucalyptus* and *Banksia* woodland with a dense understorey about one metre in height but has been observed to forage in burnt areas (Arnold *et al.*, 1991; Bamford and Bamford 1999). The preference for dense long-unburnt understorey vegetation appears to be related to shelter and refuge requirements. On the GGS this vegetation attributes are associated with wetlands and damplands of both the Spearwood and Bassendean soils. The species is thus susceptible to declining groundwater levels that lead to changes in vegetation structure surrounding wetlands and damplands.

Susceptibility of mammals to declining rainfall

There is evidence from the current scientific literature that seven mammal species considered to be extant on the GGS are likely to be impacted by declining rainfall (see studies listed in Table 5). For example, *H. chrysogaster* may be indirectly affected by low water levels in wetlands and lakes due to low rainfall levels. High and low population numbers of *R. fuscipes* are related to high and low levels of rainfall.

Declines in population numbers of *M. fuliginosus* are commonly recorded when droughts result in food shortages, for example in Western Australia, New South Wales, South Australia and southern Queensland. A study of a population in Western Australia (Arnold and Steven, 1988) found that the number of *M. fuliginosus* declined by 40% (from 800 to 500) during a drought year and remained at the reduced level for at least the next 3 years.

Reduced rainfall is also likely to have an indirect negative impact on *Macropus irma*, due to the potential reduction in dense vegetation understorey of wetlands and damplands that the species is dependent upon.

Relationship of *T. rostratus* to declining rainfall

We modelled possible changes in abundance of *T. rostratus* with incremental declines in rainfall (Figure 4). The relationship between *T. rostratus* capture rate and rainfall in the GGS was obtained using data from Wooller *et al.* (1998) (linear equation: *T. rostratus* capture rate (per 1000 trap night) = $0.119 \times \text{annual rainfall (mm)} + 2.84$; $r^2 = 0.65$) and then applied to our study area using rainfall data from the GGS. The extreme scenario modelled a 60% decline in rainfall, and resulted in a 58% decline in predicted capture rates. The annual rainfall of recent years (2006 – 2010) indicates that extremely dry years are currently occurring on the GGS, and that *T. rostratus* abundance would be predicted to be comparatively quite low (e.g. prediction of ~ 37% reduction in capture rates in 2006).

Discussion

Historic occurrences and declines of mammals

The decline of mammals on the GGS following European settlement has been dramatic with a loss of species and declines in distribution. In addition, a number of remaining species are now most likely restricted to isolated remnants. Surprisingly, few studies on mammal persistence have been conducted in the GGS area. Once species-rich, with up to 28 terrestrial native mammal species, the northern SCP has suffered a drastic decline in mammal species, with only 9 species recorded (captured or observed) in the current study, and two species (*C. concinnus* and *D. geoffroii*) considered to exist in the study area based on recent sightings outside this study. Mammal capture rates were very low, with the exception of *M. musculus* and *T.*

443 *rostratus*, which were the most frequently captured mammal species. The remaining
444 17 terrestrial mammal species recorded prior to the 1970s are considered unlikely to
445 still occur in the GGS area.

446
447 For some species, records have been sporadic throughout the last few decades. For
448 example, Ash-grey mice (*Pseudomys albocinereus*) were recorded historically on the
449 northern SCP and were trapped at a range of sites in woodland and heath habitats by
450 Kitchener *et al.* (1978). It has not been recorded since 1987, but has been recorded
451 nearby to the north-east of the study area on the Dandaragan plateau (Bamford, 1986;
452 Burbidge *et al.*, 1996) and may possibly persist in isolated pockets of habitat in the
453 GGS area (R. Davis, *pers. comm.*). Species, such as *C. concinnus*, were not recorded
454 by Kitchener *et al.* (1978) or in the current study, although recent observations of *C.*
455 *concinnus* (R. Davis, *pers. comm.*) indicate that remnant populations may exist.
456 During the current study a juvenile of an unidentified *Sminthopsis* species was
457 captured, indicating that at least one *Sminthopsis* taxon is still extant on the GGS,
458 albeit in small numbers.

459
460 A number of threatening processes have been implicated in the decline of mammal
461 fauna in the study area following European settlement. These include habitat
462 clearance and fragmentation as a result of agriculture and urbanisation (Kitchener *et*
463 *al.*, 1980; How and Dell, 2000), changed fire regimes (How *et al.*, 1987; Wilson *et al.*,
464 2010), predation by introduced foxes and cats (Kinnear *et al.*, 2002;) and possibly
465 wildlife disease (Abbott, 2008). Recently, the impacts of the plant pathogen
466 *Phytophthora cinnamomi* dieback on flora and ecosystems have been identified as
467 threatening mammal fauna on the GGS (Wilson *et al.*, 2009). Mammal declines are
468 unlikely to be attributed to a single factor, and it is more likely that each species has
469 been affected differently by various combinations of factors. Importantly, individual
470 threatening processes may have synergistic effects, resulting in compounded impacts
471 (McKenzie *et al.*, 2007; Wilson and Valentine, 2009).

472 473 **Susceptibility of mammals to declining rainfall and groundwater levels**

474 In addition to the threatening processes discussed above, declining rainfall and
475 groundwater levels due to climate change have recently been identified as major
476 threats to terrestrial mammal species on the GGS (Wilson and Valentine, 2009; Isaac

et al., 2009). There is a strong geographical pattern of the highest species loss in arid regions (McKenzie *et al.*, 2007), and recent modelling suggests that mammals in the critical weight range (i.e. non-flying mammals with an average body weight in the range 35-4200 g) in the GGS area may be very susceptible to the impacts of declining rainfall including *H. chrysogaster*, *I. obesulus* and *R. fuscipes*. Similar results have been found by Isaac *et al.* (2009), although this was in Australia's wet tropics. There is recent evidence that factors such as drought have had significant impacts on some native small mammals in southern Australian communities resulting in reduced body weight, survival and population size (e.g. Sale *et al.*, 2008; Magnusdottir *et al.*, 2008; Recher *et al.*, 2009).

In particular, the survival of *H. chrysogaster* on the GGS is critically linked to declining rainfall and groundwater levels and the persistence of wetland ecosystems. While large populations of *H. chrysogaster* have been recorded in eastern Australia (e.g. Murray Goulburn valley, Victoria, Griffith New South Wales (McNally, 1960; Woollard *et al.*, 1978) the populations on the northern SCP appear to be smaller than populations in Victoria and New South Wales. *Hydromys chrysogaster* has already suffered a significant decline in south-west Australia and is highly susceptible to loss of habitat through the contraction and drying out of lakes, the filling in and draining of wetland ecosystems for alternative land use (Burbidge and McKenzie, 1989; Lee, 1995; Morris, 2000) and loss of wetland-associated prey (Atkinson *et al.*, 2008). It is likely that food resources and low availability of suitable water bodies limits these populations.

In this study the captures at Lake Goollellal and Loch McNess indicated that there are likely to be small populations at these lakes whereas only one individual was captured at Lake Joondalup, which is a substantially larger lake. The vegetation surrounding Lake Joondalup had recently been burnt (< 3 years since last fire) and it is possible that the reduced vegetation structure and cover had impacted on the *H. chrysogaster*. In addition, unlike Lake Goollellal and Loch McNess, Lake Joondalup lacks islands located close to the lake edge, which are likely to provide feeding areas and refugia from predation or disturbance from humans.

A more recent study than the current study assessed the occurrence of *H. chrysogaster* at 39 sites in the greater Perth area (Smart, 2009). Animals were captured at only two sites, while indirect evidence of their presence (footprints, scats, midden feeding) was recorded at a further four sites. Re-sampling of the current study's sites by Smart (2009) resulted in few captures of *H. chrysogaster* (Loch McNess: n = 0; Lake Goolellal: n = 2; Lake Joondalup: n = 1), while camera trapping at Loch McNess has recently recorded two individuals (K. Bettink *pers comm*, 2011). These findings support our results that *H. chrysogaster* population numbers on the GGS are very low, and provide evidence that there may be differences in trappability or annual or seasonal differences in population numbers.

Isodon obesulus prefers scrub habitats or areas that provide dense ground cover throughout Australia (Hocking, 1990; Paull, 1995), and in Western Australia *I. obesulus* occurs in two distinct habitats across its range, open forest and dense vegetation around swamps and watercourses (Cooper, 2000a; Cooper, 2000b). The distribution of *I. obesulus* on the GGS appears to be strongly linked to dense wetland-associated vegetation around swamps and watercourses (Bamford and Bamford, 1994; Cooper, 2000a; Cooper, 2000b). Any decline in wetland vegetation associated with rainfall and groundwater decline is predicted to severely impact this species.

In eastern Australia, *R. fuscipes* is an omnivore and is considered to be a habitat generalist occurring in a wide range of habitats including coastal heath, rainforest, wet and dry forests, and woodlands (Newsome and Catling, 1979; Moro, 1991; Lunney 2008). Although, *R. fuscipes* exhibits a preference for increased vegetation cover at all structural levels, it is associated with some floristic factors (Laidlaw and Wilson, 2006). On the GGS it appears that the level of vegetation complexity and cover is only provided in vegetation surrounding wetlands, and that *R. fuscipes* will therefore be susceptible to changes in vegetation structure induced by declining rainfall and groundwater levels.

Species such as *T. rostratus* are not considered to have a high reliance upon groundwater levels as they mostly occur in upland *Banksia* woodlands. However, declining groundwater levels and reduced rainfall rates are predicted to alter these vegetation communities (Groom *et al.*, 2000) and changes in vegetation cover,

floristics composition and productivity have already been recorded (Sommer and Froend 2011). For nectar-feeding species such as *T. rostratus*, declining rainfall and a lowering groundwater table may affect the flowering period of the species upon which these marsupials feed, such as banksias. Groom *et al.* (2000) found that three species of banksia, in particular *Banksia ilicifolia* were severely impacted by drought stress within the GGS area. This was attributed to a significant lowering of the groundwater table caused by the cumulative effects of groundwater abstraction and below average rainfall (Groom *et al.*, 2000). Deaths of banksia trees on the Scott River plains (south coast of Western Australia) may also be due to a lowered ground water table, and are thought to be the cause of a corresponding decrease in honey possum populations (Phillips *et al.*, 2004). The GGS area is already experiencing declines in annual rainfall, with a 9% decline in rainfall from 1976 to 2007, and a 13% decline in rainfall from 1997 to 2007 (Government of Western Australia, 2009). As *T. rostratus* typically only lives for 1 to 3 years, there is the potential for successive dry years to affect the persistence of *T. rostratus* populations.

The loss of surface water and aquatic habitat for mammals such as *H. chrysogaster*, is likely to be catastrophic and the impacts could be quick. There is recent evidence that some lakes (e.g. Loch McNess) have undergone a sudden transition since 2006 and are now disconnected from the aquifer (DOW, 2011). This has resulted in the lowest peak water levels ever recorded and the drying out of 80% of lakes, such as Loch McNess, in summer. The disappearance of surface water would most certainly have a negative effect on the health of wetland vegetation, and thus mammal species that rely on wetland vegetation.

Management implications for mammals in a drying climate

Effective management will be needed to protect key mammal species on the GGS that are predicted to be impacted by declining rainfall and groundwater levels. Although it is impossible to control rainfall, it is possible that the management of some of the other threatening processes on the GGS (habitat loss and fragmentation, altered fire regimes and introduced predators) may increase the resilience of species such as *I. obesulus*, *H. chrysogaster* and *T. rostratus* to the threats of declining rainfall and groundwater levels. Management procedures to increase resilience could include actions such as provision of suitable refugia, establishment of ecological linkages and

baiting of predators. In addition, on the GGS a number of management interventions that aim to prevent loss of, or permanent change in, groundwater-dependent wetland systems have been employed, including turning off nearby extraction bores to reduce drawdown effects, and using artificial water supplementation pumped from the superficial aquifer (Groom *et al.* 2000; DOW 2008).

Identification and management of wetland refugia

Providing key refuge sites can buffer species from a number of threats, such as impacts of climate change. While approximately 75% of the wetlands of the SCP have been filled in or drained (Commonwealth of Australia, 1997), groundwater modelling predicts that those wetlands closest to the top of the Gnangara aquifer located in the northeast (e.g. Yeal Nature Reserve) will be significantly affected by lower water levels and terrestrialisation in the future, up to an 11 metre decline by 2030 (Government of Western Australia, 2009). Smaller water level declines are likely by 2030 for Lakes Joondalup and Goollelal, the most southern of the long chain coastal wetlands. A preliminary set of wetlands including both permanently and seasonally inundated wetlands (i.e. lakes and sumplands) have been identified on the GGS as significant in terms of biodiversity values including wetland associated mammals and ecological function (Government of Western Australia, 2009; Horwitz *et al.*, 2009). It is recommended that key refugia be selected from the identified wetlands and managed so as to protect them from further loss and modification, particularly along lake banks where buffers need to be implemented. Habitat requirements that need to be maintained to encourage successful breeding and feeding of *H. chrysogaster* include areas suitable for location of burrows and nests such as steep banks, the presence of logs, rocks or reeds as feeding areas, and adequate riparian and stream vegetation cover (Woollard *et al.*, 1978; Smart, 2009).

Of major importance for the management of wetland refugia will be the implementation of specific ecological fire regimes to protect them from inappropriate fires, such as frequent lethal and intense fires (Horwitz *et al.*, 2009; Wilson *et al.*, 2010). In particular the ecological fire regimes must ensure the retention and protection of the long-unburnt wetland-associated vegetation (Wilson *et al.*, 2010).

Predator control is likely to be an essential component of management for the wetland refugia. In the GGS area *I. obesulus* is restricted to dense wetland-associated vegetation, although it occupies upland habitat in areas where predators have been suppressed (Bamford and Bamford, 1994; Valentine *et al.*, 2009). This was interpreted as a response to the diminution of the threat of predation (Bamford and Bamford, 1994). Although foxes have been recorded regularly in fauna surveys on the northern SCP (Kitchener *et al.*, 1978; Valentine *et al.*, 2009), and there is evidence that both *I. obesulus* and *M. irma* have increased in numbers since baiting commenced (C. Rafferty *pers. comm.* 2009), there is no coordinated baiting program throughout the study area.

Identification and management of ecological linkages

A major recommendation of the Gnangara Sustainability Strategy (Government of Western Australia, 2009) is to establish ecological linkages covering 9 000 ha across the GGS. The ecological linkages proposed include viable remnants (e.g. conservation reserves, Bush Forever sites), vegetated waterways and drainage lines, large remnant vegetation patches within pine plantations, and areas of pine plantations that will be rehabilitated (9 000 ha post-harvest). It is vital that these linkages be managed to support fauna species, such as *H. chrysogaster* and *T. rostratus*.

During the current study *T. rostratus* was recorded both in native vegetation and within small vegetation remnants in the pine plantations across the GGS. The ecological linkages need to be designed and managed to provide food resources and vegetation of suitable post-fire age and structure for the species. Although *T. rostratus* in south-west Western Australia is known to return to burnt areas within 2 – 4 years since fire (Bamford, 1986; Richardson and Wooller, 1991; Everaardt, 2003), higher densities are recorded in older vegetation, with peaks in abundance in vegetation 20 – 30 years since last burnt (Everaardt, 2003; Bradshaw *et al.*, 2007). It is recommended that native remnant vegetation within the ecological linkages consist predominantly of preferred post fire age for this species (20-30 years since last fire).

There is little information on the distribution, population dynamics and habitat requirements of key mammal species on the GGS that are predicted to be severely impacted by declining rainfall and groundwater levels. The population dynamics and

breeding of *H. chrysogaster* in the wetlands within the GGS, in particular what the population sizes are, how the populations fluctuate seasonally or annually, and the stability of the populations needs to be determined. This is a species that will require substantial consideration and careful management as any reduction in the viability of wetlands and food resources on the GGS as a result of declining rainfall and groundwater are likely to significantly impact on breeding and feeding of water rats. Species that exhibit a positive relationship between population abundance and rainfall, such as *T. rostratus*, are likely to decline under future rainfall scenarios. As a relatively short-lived species, *T. rostratus* may also be susceptible to extreme weather events, such as sequential very low rainfall years. Without detailed biological and ecological information on the species of the GGS area, it is difficult to make accurate predictions on the extent of the impacts of declining rainfall. The fauna survey sites that were established in the GGS area for this project should provide a basis for monitoring programs for these species.

Acknowledgements

This work was funded by the Gngangara Sustainability Strategy, Government of Western Australia, and we thank in particular Dr. Melanie Strawbridge, Department of Water, WA. We also thank the other members of the Department of Environment and Conservation, WA – Gngangara Sustainability Strategy team: Project manager Paul Brown and team members Dr Mark Garkaklis, Janine Kinloch, Marnie Swinburn, Tracy Sonneman, Natalia Huang, Brent Johnson, David Mickle, Katie Montgomery, Janine Kuehs and Lara Ryniker. Thank you also to the many field volunteers and the Department of Environment and Conservation Swan Coastal District staff for their assistance during this project. Our animal ethics permits were: AEC Approval Number: DEC AEC 37 / 2007, and the Department of Environment and Conservation Licence to Take Fauna for Scientific Purposes Number: SC 000826.

References

Abbott, I. (2008) Historical perspectives of the ecology of some conspicuous vertebrate species in south-west Western Australia. *Conservation Science Western Australia*, 6, 1-214.

- 678 Arnold, G. W., Smith, G. T. & Brooker, M. G. (1991) Whiteman Park fauna survey:
679 final report. Prepared for the State Planning Commission, Whiteman Park,
680 Western Australia. CSIRO Division of Wildlife and Ecology.
- 681 Arnold, G. W. & Steven, D. E. (1988) Variations in Distribution of Western Grey
682 Kangaroos, *Macropus fuliginosus ocydromus*, in the Tutanning Nature
683 Reserve and their Impact on Adjacent Farmland. *Australian Wildlife*
684 *Research*, 15, 119-128.
- 685 Abensperg-Traun, M. (1994) The influence of climate on patterns of termite eating in
686 Australian mammals and lizards. *Australian Journal of Ecology*, 19, 65-71.
- 687 Atkinson, C. A., Lund, M. A. & Morris, K. D. (2008) Biblio Rakali: the Australian
688 water rat, *Hydromys chrysogaster* Geoffroy, 1804 (Muridae:
689 Hydromyinae), a subject-specific bibliography. *Conservation Science*
690 *Western Australia*, 7, 65-71.
- 691 Bamford, M. J. (1986) The dynamics of small vertebrates in relation to fire in *Banksia*
692 woodland near Perth, Western Australia. PhD Thesis, Murdoch
693 University, Perth, Western Australia.
- 694 Bamford, M. J. & Bamford, A. R. (1990) *Yellagonga Regional Park: a preliminary*
695 *survey of vertebrate fauna*. Department of Conservation and Land
696 Management, Perth, Western Australia.
- 697 Bamford, M. J. & Bamford, A. R. (1994) Survey of the Quenda *Isodon obesulus* and
698 other medium-sized, native mammals, December 1993. Department of
699 Planning and Urban Development, Perth, Western Australia.
- 700 Bamford, M. J. & Bamford, A. R. (1999) The estimation of the number of Western
701 Grey Kangaroos in Whiteman Park; May-June 1999. Whiteman Park
702 Authority, and Department of Planning and Urban Development, Perth,
703 Western Australia.
- 704 Bamford, M. J. & Bamford, A. R. (2003) Vertebrate Fauna; Values and Groundwater
705 Dependence in the Gnangara Study Area. Report for the Water and Rivers
706 Commission, Perth, Western Australia.
- 707 Beard, J. S. (1984) Biogeography of the Kwongan. In: Pate, J. S. & Beard, J. S. (Eds.)
708 *Kwongan. Plant life of the Sandplain*. University of Western Australia
709 Press, Nedlands, Western Australia.
- 710 Bradshaw, S. D., Phillips, R. D., Tomlinson, S., Holley, R. J., Jennings, S. &
711 Bradshaw, F. J. (2007) Ecology of the Honey Possum, *Tarsipes rostratus*,

- 712 in Scott National Park, Western Australia. *Australian Mammalogy*, 29, 25-
713 38.
- 714 Braithwaite, R. W. & Muller, W. J. (1997) Rainfall, groundwater and refuges:
715 Predicting extinctions of Australian tropical mammal species. *Australian*
716 *Journal of Ecology*, 22, 57-67.
- 717 Bryant, K. A. (2004) The mating system and reproduction in the honey possum,
718 *Tarsipes rostratus*: a life-history and genetical perspective, PhD Thesis.
719 *School of Biological Sciences and Biotechnology*. Murdoch University,
720 Perth, Western Australia.
- 721 Burbidge, A. A. & McKenzie, N. L. (1989) Patterns in the modern decline of Western
722 Australia's vertebrate fauna: causes and conservation implications.
723 *Biological Conservation*, 50, 143-198.
- 724 Burbidge, A. H., Boscacci, L. J., Alford, J. J. & Keighery, G. J. (1996) A biological
725 survey of Boonanarring nature reserve. *CALMScience*, 2, 153-187.
- 726 Cairns, S. C., Grigg, G. C., Beard, L. A., Pople, A. R. & Alexander, P. (2000)
727 Western grey kangaroos, *Macropus fuliginosus*, in the South Australian
728 pastoral zone: populations at the edge of their range. *Wildlife Research*,
729 27, 309-318.
- 730 Carthew, S. M., Cadzow, B. R. & Foulkes, J. N. (2008) Western Pygmy Possum. In:
731 Van Dyck, S. M. & Strahan, R. (Eds.) *The Mammals of Australia - 3rd*
732 *Edition*. Reed New Holland Publishers, Sydney, Australia.
- 733 Caughley, G. (1987) Ecological relationships. In: Caughley, G., Shepherd, N. & Short,
734 J. (Eds.) *Kangaroos: Their Ecology and Management in the Sheep*
735 *Rangelands of Australia*. Pp. 119-134. Cambridge University Press,
736 Cambridge.
- 737 Caughley, G., Grigg, G. C. & Smith, L. (1985) The effect of drought on kangaroo
738 populations. *Journal of Wildlife Management*, 49, 679-685.
- 739 Caughley, J., Bayliss, P. & Giles, J. (1984) Trends in kangaroo populations in western
740 New South Wales and their relation to rainfall. *Australian Wildlife*
741 *Research*, 11, 415-422.
- 742 Commonwealth of Australia (1997) Nationally agreed criteria for the establishment of
743 a comprehensive, adequate and representative reserve system for forests in
744 Australia. ANZECC/MCFFA National Forest Policy Statement
745 Implementation Sub-Committee, Canberra, Australia.

- 746 Cooper, M. L. (1998) Geographic variation in size and shape in the southern brown
 747 bandicoot, *Isoodon obesulus* (Peramelidae : Marsupialia), in Western
 748 Australia. *Australian Journal of Zoology*, 46, 145-152.
- 749 Cooper, M. L. (2000a) Random amplified polymorphic DNA analysis of southern
 750 brown bandicoot (*Isoodon obesulus*) populations in Western Australia
 751 reveals genetic differentiation related to environmental variables.
 752 *Molecular Ecology*, 9, 469-479.
- 753 Cooper, M. L. (2000b) Temporal variation in skull size and shape in the southern
 754 brown bandicoot, *Isoodon obesulus* (Peramelidae : Marsupialia) in
 755 Western Australia. *Australian Journal of Zoology*, 48, 47-57.
- 756 Coulson, G. (2008) Western Grey Kangaroo (*Macropus fuliginosus*). In: Van Dyck, S.
 757 M. & Strahan, R. (Eds.) *The Mammals of Australia - 3rd Edition.*, Reed
 758 New Holland Publishers, Sydney, Australia.
- 759 DOW (2008) Gnangara groundwater areas - Water Management Plan (draft for public
 760 comment). Department of Water, Perth, Western Australia.
- 761 DOW (2011) *Perth shallow groundwater systems investigation: Loch McNess*,
 762 Department of Water, Hydrogeological record series report no. HG43.
 763 Government of Western Australia, Perth, Western Australia.
- 764 Drew, M. (1998) Patterns in terrestrial vertebrate fauna populations in Banksia
 765 woodlands of Whiteman Park. Murdoch University, Perth, Australia.
- 766 Eamus, D., Hatton, T., Cook, P. & Colvin, C. (2006) *Ecohydrology: Vegetation*
 767 *function, water and resource management*. CSIRO Publishing,
 768 Melbourne, Australia.
- 769 Ecologia Environmental Consultants (1990) Southern brown bandicoot *Isoodon*
 770 *obesulus* Biological assessment survey. City of Mandurah Town Planning,
 771 Mandurah, Western Australia.
- 772 Ecologia Environmental Consultants (1997) Mitchell Freeway: Romeo Road to Perth-
 773 Lancelin Road: biological assessment. Main Roads Western Australia.
- 774 Everaardt, A. (2003) The impact of fire on the Honey Possum, *Tarsipes rostratus*, in
 775 the Fitzgerald River National Park, Western Australia (PhD Thesis).
 776 Murdoch University, Perth, Western Australia.
- 777 Friend, A. (1996) The Department of Conservation and Land Management Bandicoot
 778 Survey 1991 - 1996. *Database of Bandicoot sightings reported by the*

- 779 *public*. Department of Conservation and Land Management, Como,
780 Western Australia.
- 781 Government of Western Australia (2000) Bush Forever Volume 2: Directory of Bush
782 Forever Sites. Department of Environmental Protection, Perth, Australia.
- 783 Government of Western Australia (2009) Draft Gnamptogaster Sustainability Strategy.
784 Department of Water and the Gnamptogaster Sustainability Strategy, Perth,
785 Western Australia.
- 786 Groom, P. K., Froend, R. H. & Mattiske, E. M. (2000) Impact of groundwater
787 abstraction on a Banksia woodland, Swan Coastal Plain, Western
788 Australia. *Ecological Management & Restoration*, 1, 117-124.
- 789 Heddle, E. M., Loneragan, D. W. & Havel, J. J. (1980) Vegetation complexes of the
790 Darling System, Western Australia. *Atlas of natural resources, Darling*
791 *System, Western Australia*. Department of Conservation and Environment,
792 Perth, Western Australia.
- 793 Hocking, G. (1990) Status of bandicoots in Tasmania. In: Seebeck, J., Brown, P.,
794 Wallis, R. & Kemper, C. (Eds.) *Bandicoots and Bilbies*. Surrey Beatty and
795 Sons, Chipping North, New South Wales, Australia.
- 796 Horwitz, P., Sommer, B. & Hewitt, P. (2009) Chapter Five: Wetlands - Changes,
797 Losses and Gains. In: Wilson, B. A. & Valentine, L. E. (Eds.) *Biodiversity*
798 *values and threatening processes of the Gnamptogaster groundwater system*. A
799 report to the Gnamptogaster Sustainability Strategy and the Department of
800 Environment and Conservation, Perth, Western Australia.
- 801 How, R. A. & Dell, J. (1993) Vertebrate Fauna of the Perth metropolitan region:
802 consequences of a modified environment. In: Hipkins, M. (Ed.) *Urban*
803 *Bush Management*. Australian Institute of Urban Studies, Western
804 Australia.
- 805 How, R. A. & Dell, J. (1994) The zoogeographic significance of urban bushland
806 remnants to reptiles in the Perth region, Western Australia. *Pacific*
807 *Conservation Biology*, 1, 132-140.
- 808 How, R. A. & Dell, J. (2000) Ground vertebrate fauna of Perth's vegetation remnants:
809 Impact of 170 years of urbanization. *Pacific Conservation Biology*, 6, 198-
810 217.

- 811 How, R. A., Harvey, M. S., Dell, J. & Waldock, J. M. (1996) Ground Fauna of Urban
812 Bushland Remnants in Perth: Report to the Australian Heritage
813 Commission.
- 814 How, R. A., Dell, J. & Humphreys, W. F. (1987) The ground vertebrate fauna of
815 coastal areas between Busselton and Albany, Western Australia. *Records*
816 *of the Western Australian Museum*, 13, 423-443.
- 817 Hughes, L. (2000) Biological consequences of global warming: is the signal already
818 apparent? *Trends in Ecology & Evolution*, 15, 56-61.
- 819 IOCI (2005) How our rainfall has changed in the South West. Indian Ocean Climate
820 Initiative (IOCI), Department of Environment, Perth, Western Australia.
- 821 IPCC (2007) Climate Change 2007: impacts, adaptation and vulnerability.
822 Contribution of working group II to the fourth assessment report of the
823 IPCC. Intergovernmental Panel on Climate Change, Geneva.
- 824 Isaac, J. L., Vanderwal, J., Williams, S. E. & Johnson, C. N. (2009) Resistance and
825 resilience: quantifying relative extinction risk in a diverse assemblage of
826 Australian tropical rainforest vertebrates. *Diversity and Distributions*, 15,
827 280-288.
- 828 Johnson, C. N. (2006) *Australia's mammal extinctions: A 50 000 year history*.
829 Cambridge University Press, Cambridge.
- 830 Jones, D. B. & Preston, B. L. (2006) Climate change impacts, risk and the benefits of
831 mitigation. CSIRO Publishing, Australia.
- 832 Kinhill Pty Ltd (1997) Perth to Darwin National Highway, Reid Highway to Maralla
833 Road. Shire of Swan, Perth, Western Australia.
- 834 Kinnear, J. E., Sumner, N. R. & Onus, M. L. (2002) The red fox in Australia - an
835 exotic predator turned biological agent. *Biological Conservation*, 108,
836 335-359.
- 837 Kitchener, D. J., Chapman, A. R. & Barron, G. (1978) Mammals of the northern Swan
838 Coastal Plain. *Faunal Studies of the Northern Swan Coastal Plain - a*
839 *consideration of past and future changes*. Western Australian Museum,
840 Department of Conservation and Environment, Perth, Western Australia.
- 841 Kitchener, D. J., Chapman, A. R. & Muir, B. G. (1980) The Conservation Value for
842 Mammals of Reserves in the Western Australian Wheatbelt. *Biological*
843 *Conservation*, 18, 179-207.

- 844 Laidlaw, W. S. & Wilson, B. A. (2006) Habitat utilisation by small mammals in a
845 coastal heathland exhibiting symptoms of *Phytophthora cinnamomi*
846 infestation. *Wildlife Research*, 33, 639-649.
- 847 Lee, A. K. (1995) *The Action Plan for Australian rodents*. Environment Australia,
848 Canberra, Australia.
- 849 Lunney, D. (2008) Bush Rat. In: Van Dyck, S. M. & Strahan, R. (Eds.) *The Mammals*
850 *of Australia - 3rd Edition*. Reed New Holland Publishers, Sydney,
851 Australia.
- 852 Lunney, D., Cullis, B. & Eby, P. (1987) Effects of logging and fire on small mammals
853 in Mumbulla State Forest, near Bega, New South Wales. *Australian*
854 *Wildlife Research*, 14, 163-181.
- 855 Magnúsdóttir, R., Wilson, B. A. & Hersteinsson, P. (2008) Dispersal and the
856 influence of rainfall on a population of swamp antechinus (*Antechinus*
857 *minimus maritimus*). *Wildlife Research*, 35, 446-454.
- 858 Malcolm, J., Liu, C., Neilson, R., Hansen, L. & Hannah, L. (2006) Global Warming
859 and extinctions of endemic species from biodiversity hotspots.
860 *Conservation Biology*, 20, 538-548.
- 861 Maxwell, S., Burbidge, A. A. & Morris, K. (1996) Action Plan for Australian
862 Marsupials and Monotremes. Environment Australia.
- 863 McKenzie, N. L., Burbidge, A., Baynes, A., Brereton, R. N., Dickman, C. R., Gordon,
864 G., Gibson, L. A., Menkhorst, A. C., Robinson, A. C., Williams, M. R. &
865 Woinarski, J. C. Z. (2007) Analysis of factors implicated in the recent
866 decline of Australia's mammal fauna. *Journal of Biogeography*, 34, 597-
867 611.
- 868 McNally, J. (1960) The biology of the water rat *Hydromys chrysogaster* Geoffroy
869 (Muridae: Hydromyinae) in Victoria. *Australian Journal of Zoology*, 8,
870 170 - 180.
- 871 Mitchell, D., Williams, K. & Desmond, A. (2003) Swan Coastal Plain 2 (SWA 2-
872 Swan Coastal Plain Sub-region). In: May, J. E. & McKenzie, N. L. (Eds.)
873 *A biodiversity audit of Western Australia's 53 Biogeographical subregions*
874 *in 2002*. Department on Conservation and Land Management, Perth,
875 Western Australia.
- 876 Moro, D. (1991) The distribution of small mammal species in relation to heath
877 vegetation near Cape Otway, Victoria. *Wildlife Research*, 18, 593-604.

- 878 Morris, K. D. (2000) The status and conservation of native rodents in Western
879 Australia. *Wildlife Research*, 27, 405-419.
- 880 Morton, S. R. (1990) The impact of European settlement on the vertebrate animals of
881 arid Australia: a conceptual model. *Proceedings of the Ecological Society
882 of Australia*, 16, 201-213.
- 883 Nevill, J. C., Hancock, P. J., Murray, B. R., Ponder, W. F., Humphreys, W. F.,
884 Phillips, M. L. & Groom, P. K. (2010) Groundwater-dependent
885 ecosystems and the dangers of groundwater overdraft: a review and an
886 Australian perspective. *Pacific Conservation Biology*, 16, 187-208.
- 887 Newsome, A. E. & Catling, P. C. (1979) Habitat preferences of mammals inhabiting
888 heathlands of warm temperate coastal, montane and alpine regions of
889 southeastern Australia. In: Specht, R. L. (Ed.) *Ecosystems of the world,
890 vol. 91. Heathlands and related shrublands of the world*. Elsevier,
891 Amsterdam.
- 892 Olsen, P. D. (1985) Water-rat, *Hydromys chrysogaster*. In: Strahan, R. (Ed.) *The
893 Mammals of Australia*. Reed Books, Chatswood.
- 894 Parmesan, C. & Yohe, G. (2003) A globally coherent fingerprint of climate change
895 impacts across natural systems. *Nature*, 421, 37-42.
- 896 Parrott, M. L., Ward, S. J., Temple-Smith, P. D. & Selwood, L. (2007) Effects of
897 drought on weight, survival and breeding success of agile antechinus
898 (*Antechinus agilis*), dusky antechinus (*A. swainsonii*) and brush rats
899 (*Rattus fuscipes*). *Wildlife Research* 21, 1-9.
- 900 Paull, D. (1995) The distribution of the southern brown bandicoot (*Isoodon obesulus*
901 *obesulus*) in South Australia. *Wildlife Research*, 22, 585-600.
- 902 Phillips, R., S., Tomlinson, S., Jennings, B. & Holley, R. (2004) Summer
903 ecophysiology of the honey possum (*Tarsipes rostratus*) on the south
904 coast of Western Australia (Honours thesis). University of Western
905 Australia, Perth, Western Australia.
- 906 Pitman, A. J. & Perkins, S. E. (2008) Regional projections of future seasonal and
907 annual changes in rainfall and temperature over Australia based on skill-
908 selected AR(4) models. *Earth Interactions*, 12, 1-50.
- 909 Playford P. E., Cockbain A. E. & Low, G. H. (1976) Geology of the Perth Basin,
910 Western Australia. *Geological Surveillance Western Australia Bulletin*
911 124, 1-130.

- 912 Recher, H. F., Lunney, D. & Matthews, A. (2009) Small mammal populations in a
913 eucalypt forest affected by fire and drought. I. Long-term patterns in an
914 era of climate change. *Wildlife Research*, 36, 143-158.
- 915 Rhind, S. G. & Bradley, J. S. (2002) The effect of drought on body size, growth and
916 abundance of wild brush-tailed phascogales (*Phascogale tapoatafa*) in
917 south-western Australia. *Wildlife Research*, 29, 235-245.
- 918 Richardson, K. C. & Wooller, R. D. (1991) The effects of fire on Honey Possum
919 populations. Report prepared for World Wide Fund for Nature, Perth,
920 Western Australia.
- 921 Sale, M. G., Wilson, B. A. & Arnould, J. P. Y. (2008) Factors influencing population
922 dynamics in island and mainland populations of swamp atechinus
923 (*Antechinus minimus*; Marsupialia). *Australian Journal of Zoology*, 56,
924 187-194.
- 925 Seebeck, J. & Menkhorst, P. (2000) Status and conservation of the rodents of
926 Victoria. *Wildlife Research*, 27, 357-369.
- 927 Semeniuk, C. A., Semeniuk, V., Cresswell, I. D. & Marchant, N. G. (1990) Wetlands
928 of the Darling System, southwestern Australia: a descriptive classification
929 using pattern and form. *Journal of the Royal Society of Western Australia*,
930 72, 109-121.
- 931 Short, J. & Smith, A. (1994) Mammal decline and recovery in Australia. *Journal of*
932 *Mammalogy*, 75, 288-297.
- 933 Smart, C. (2009) An ecological study of the Australian water rat (*Hydromys*
934 *chrysogaster*) in the greater Perth region, Western Australia:
935 Environmental and biological factors influencing distribution. *School of*
936 *Animal Biology*. The University of Western Australia, Perth, Western
937 Australia.
- 938 Sommer, B. & Froend, R. (2011) Resilience of phreatophytic vegetation to
939 groundwater drawdown: is recovery possible under a drying climate?
940 *Journal of Ecohydrology*, 4, 67-82.
- 941 Valentine, L. E., Wilson, B. A., Reaveley, A., Huang, N., Johnson, B. & Brown, P. H.
942 (2009) *Patterns of ground-dwelling vertebrate biodiversity in the*
943 *Gnangara Sustainability Strategy study area*. Unpublished report prepared
944 by Department of Environment and Conservation for the Gnangara
945 Sustainability Strategy, Perth, Western Australia.

- 946 Vogwill, R. I. J., Mchugh, S. L., O'Boy, C. A. & Yu, X. (2008) PRAMS scenario
 947 modelling for water management of the Gnangara Groundwater Mound,
 948 HG 21. Department of Water, Perth, Western Australia.
- 949 Walther, G. R., Post, E., Convery, P., Menzel, A., Parmesan, C., Beebee, T. J. C.,
 950 Fromentin, J., Hoegh-Guldberg, O. & Bairlein, F. (2002) Ecological
 951 responses to recent climate change. *Nature* 416, 389-395.
- 952 Wilson, B. A., Bleby, K., Valentine, L. E., Swinburn, M. L. & Kuehs, J. M. (2010)
 953 Impacts of Fire on Biodiversity of the Gnangara Groundwater System.
 954 Department of Environment and Conservation, Perth, Western Australia.
- 955 Wilson, B. A., Lock, M. & Magnusdottir, R. (2007) Influence of rainfall on native
 956 mammal populations in the Eastern Otway Ranges, Southeastern Australia
 957 (abstract). In: Rokich, D., Wardell-Jones, G. & Yates, C. (Eds.) *MedEcos*
 958 *XI. The International Mediterranean Ecosystems conference, Perth,*
 959 *Western Australia.* Kings Park and Botanic Garden, Perth Western
 960 Australia.
- 961 Wilson, B. A., Kinloch, J., Sonneman, T. & Swinburn, M. (2009) Chapter Nine:
 962 Distribution and impacts of *Phytophthora cinnamomi*. In: Wilson, B. A. &
 963 Valentine, L. E. (Eds.) *Biodiversity values and threatening processes of*
 964 *the Gnangara groundwater system.* A report to the Gnangara
 965 Sustainability Strategy and the Department of Environment and
 966 Conservation. Perth, Western Australia.
- 967 Wilson, B. A. & Valentine, L. E. (Eds.) (2009) *Biodiversity values and threatening*
 968 *processes of the Gnangara groundwater system.* Department of
 969 Environment and Conservation, Perth, Western Australia.
- 970 Woinarksi, J. C. Z., Armstrong, M., Brennan, K., Fisher, A., Griffiths, A. D., Hill, B.,
 971 Milne, D. J., Palmer, C., Ward, S., Watson, M., Winderlich, S. and Young,
 972 S. (2010) Monitoring indicates rapid and severe decline of native small
 973 mammals in Kakadu National Park, northern Australia. *Wildlife Research*,
 974 37, 116-126.
- 975 Woinarski, J. C. Z., Milne, D. J. & Wanganeen, G. (2001) Changes in mammal
 976 populations in relatively intact landscapes of Kakadu National Park,
 977 Northern Territory, Australia. *Austral Ecology*, 26, 360-370.

- 978 Woollard, P., Vestjens, W. J. M. & Maclean, L. (1978) The ecology of the Eastern
979 Water Rat *Hydromys chrysogaster* at Griffith, N.S.W.: food and feeding
980 habits. *Australian Wildlife Research*, 5, 59 - 73.
- 981 Wooller, R. D., Renfree, M. B., Russell, E. M., Dunning, A., Green, S. W. & Duncan,
982 P. (1981) Seasonal changes in a population of the nectar feeding marsupial
983 *Tarsipes spencerae* (Marsupialia: Tarsipedidae). *Journal of Zoology*
984 (London), 195, 267-279.
- 985 Wooller, R. D., Richardson, K. C., Garavanta, C. A. M., Saffer, V. M., Anthony, C. &
986 Wooller, S. J. (1998) The influence of annual rainfall upon capture rates of
987 a nectar-dependent marsupial. *Wildlife Research*, 25, 165-169.
- 988 Yesertener, C. (2007) Assessment of the declining groundwater levels in the
989 Gngangara groundwater mound, Western Australia. *Hydrogeological*
990 *Record Series HG14*. Department of Water, Perth, Western Australia.
- 991
992
993

994 Table 1: Historic occurrence of non-volant native mammal species in the GGS and the
 995 time periods in which they have been recorded (Arnold *et al.*, 1991; Bamford, 1986;
 996 1990; 1994; 1999; Burbidge *et al.*, 1996; Drew, 1998; Ecologia Environmental
 997 Consultants, 1990; 1997; Friend, 1996; How and Dell, 2000; How *et al.*, 1996; Kinhill
 998 Pty Ltd, 1997; Kitchener *et al.*, 1978). Endemicity: RE (regional SWWA), WA
 999 (restricted to WA). Grey highlight refers to mammals considered to be extant in the
 1000 GGS.

Species	Common name	Endemicity	1800-1850	1850-1900	1900-1950	1950-1977	1977-2000	2000-2008
<i>Macropus fuliginosus</i>	Western grey kangaroo	AUS	✓	✓	✓	✓	✓	✓
<i>Macropus irma</i>	Western brush wallaby * (P4)	RE	✓	✓	✓	✓	✓	✓
<i>Macropus eugenii</i>	Tammar wallaby	AUS		✓	✓		✓	
<i>Petrogale lateralis lateralis</i>	Black-footed rock-wallaby* (V,S1)	WA		✓				
<i>Onychogalea lunata</i>	Crescent nailtail wallaby* (E,S2)	AUS		✓		✓ **		
<i>Bettongia penicillata ogilbyi</i>	Woylie or brushtail bettong* (S1)	AUS	✓	✓	✓			
<i>Bettongia lesueur lesueur</i>	Burrowing bettong or boodie* (V,S1)	AUS	✓	✓		✓ **		
<i>Setonix brachyurus</i>	Quokka* (V,S1)	RE	✓		✓	✓		
<i>Lagostrophus fasciatus fasciatus</i>	Banded hare-wallaby* (V,S1)	WA		✓				
<i>Trichosurus vulpecula</i>	Brushtail possum	AUS			✓	✓	✓	✓
<i>Pseudocheirus occidentalis</i>	Western ringtail possum* (V,S1)	RE	✓		✓	✓		
<i>Cercartetus concinnus</i>	Western pygmy possum	AUS		✓	✓	✓		✓
<i>Tarsipes rostratus</i>	Honey possum	RE		✓	✓	✓	✓	✓
<i>Isodon obesulus fusciventer</i>	Southern brown bandicoot or quenda* (P5)	WA			✓	✓	✓	✓
<i>Macrotis lagotis</i>	Greater bilby* (S1)	AUS		✓	✓			
<i>Dasyurus geoffroii</i>	Chuditch* (V,S1)	AUS		✓	✓	✓	✓	✓
<i>Antechinus flavipes leucogaster</i>	Mardo	WA	✓	✓	✓			
<i>Parantechinus apicalis</i>	Dibbler* (V,S1)	RE				✓ **		
<i>Phascogale tapoatafa</i>	Brush-tailed phascogale* (S1)	AUS	✓					
<i>Sminthopsis crassicaudata</i>	Fat-tailed dunnart	AUS			✓	✓		
<i>Sminthopsis griseoventer/ Sminthopsis dolichura †</i>	Grey-bellied dunnart/Little long-tailed dunnart	AUS	✓		✓			✓***
<i>Sminthopsis granulipes</i>	White-tailed dunnart	WA			✓			
<i>Myrmecobius fasciatus</i>	Numbat* (V,S1)	AUS	✓		✓			
<i>Rattus fuscipes</i>	Southern bush rat	AUS				✓	✓	✓
<i>Rattus tunneyi</i>	Tunney's rat	AUS				✓ ?		
<i>Pseudomys albocinereus</i>	Ash grey mouse or noodji	WA	✓			✓	✓	
<i>Hydromys chrysogaster</i>	Water rat or rakali* (P4)	AUS			✓	✓	✓	✓
<i>Tachyglossus aculeatus</i>	Echidna	AUS			✓	✓	✓	✓

1001 * Threatened species, threat status is in brackets: IUCN Red List and EPBC Act: 'E' – Endangered

1002 (species considered to be facing a very high risk of extinction in the wild) and 'V' – Vulnerable

1003 (species considered to be facing a high risk of extinction in the wild); WA Wildlife Conservation Act

1004 1950: 'S1' – Schedule 1 (fauna that is rare or likely to become extinct and 'S2' – Schedule 2 (presumed
1005 extinct fauna); and Department of Environment and Conservation Priority Fauna List (i.e. taxa that
1006 have not yet been adequately surveyed to be listed under Schedule 1 or 2: 'P4' – Priority 4 (taxa which
1007 are considered to have been adequately surveyed, or for which sufficient knowledge is available, and
1008 which are considered not currently threatened or in need of special protection, but could be if present
1009 circumstances change. These taxa are usually represented on conservation lands); and 'P5' – Priority 5
1010 (taxa which are not considered threatened but are subject to a specific conservation program, the
1011 cessation of which would result in the species becoming threatened within five years). ** superficial
1012 cave deposit. † *Sminthopsis griseoventer* and *S. dolichura* were previously part of the of *S. murina*
1013 complex and during the WAM 1977-78 survey were recorded as such. They are now separate species
1014 whose distribution includes the NSCP. The distribution of *S. murina* is now restricted to Eastern
1015 Australia. *** One *Sminthopsis sp.* individual was trapped during the GSS fauna survey. It is likely to
1016 have been *Sminthopsis griseoventer*, which has been recorded recently not far from this location. ?
1017 indicates uncertainty regarding species record.
1018

1019 Table 2: Species captured during pit-fall trapping or age and Elliott trapping during
 1020 general fauna surveys throughout the GGS study area, excluding targeted fauna
 1021 surveys.

			No. sites	Landform unit ^a			Vegetation type ^b				
	Pit-fall traps	Cage or Elliott traps	Total (40)	Q (2)	Sp (20)	Bn (18)	Cs (2)	Tt (4)	Jh (4)	Me (14)	Bk (16)
Species captured	Abundance			Presence / Absence			Presence / Absence				
<i>Tachyglossus aculeatus</i>	0	3	3	0	0	1	0	0	0	1	1
<i>Sminthopsis sp</i>	1	0	1	0	0	1	0	0	0	0	1
<i>Tarsipes rostratus</i>	26	0	14	0	1	1	0	0	1	1	1
<i>Mus musculus</i>	77	36	27	1	1	1	1	1	1	1	1
<i>Rattus fuscipes</i>	1	1	2	0	1	0	0	0	0	1	0
<i>Rattus rattus</i> *	0	2	2	0	1	0	0	1	0	1	0
Species Richness	4	4	NA	1	4	4	1	2	2	5	4
Total individuals	105	42	NA	21	88	38	21	11	11	49	55

^a Landform Unit: Q = Quindalup, SP = Spearwood Dunes, Bn = Bassendean Dunes

^b Vegetation Type: Cs = Coastal Scrub, Tt = tuart forest, Jh = jarrah forest, Me = *Melaleuca* wetlands, Bk = *Banksia* woodlands

1025 Table 3: Captures per 100 trap nights of mammal species captured at sites targeted for trapping *Isoodon obesulus fusciventer* and *Hydromys*
1026 *chrysogaster*.

	Muckenburra NR	Yeal NR	Lake Gnangara	Little Badgerup Swamp	Neaves Road NR	Twin Swamps NR	Maralla Road NR	Nowergup NR	Loch McNess*	Lake Joondalup**	Lake Goolellal**
<i>Isoodon obesulus fusciventer</i> Survey											
<i>I. obesulus fusciventer</i>	0.0	0.0	7.5	2.5	6.5	16.3	0.0	1.3	0.0		
<i>Rattus fuscipes</i>	1.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	25.0		
<i>Rattus rattus</i>	1.7	3.0	7.5	0.0	18.8	3.8	2.5	0.0	10.5		
<i>Mus musculus</i>	0.0	1.0	0.0	2.5	6.3	2.5	1.3	2.5	1.3		
<i>Hydromys chrysogaster</i> Survey											
<i>H. chrysogaster</i>									13.6	1.3	20.0
<i>Rattus fuscipes</i>									9.0	0.0	0.0
<i>Rattus rattus</i>									4.6	0.0	0.0
<i>Mus musculus</i>									0.0	2.5	0.0

* Located in Yanchep National Park; ** Located in Yellagonga RP

1029 Table 4: Mammal species predicted to decline in abundance or range due to a low,
 1030 moderate or high dependence on groundwater levels. Adapted from Bamford and
 1031 Bamford (2003). ('Low' = 'Low' category in Bamford and Bamford (2003), 'Strong'
 1032 = 'Moderate' and 'High' categories in Bamford and Bamford (2003), and 'High' =
 1033 'Very High' category in Bamford and Bamford (2003).
 1034

Family	Species name	Common name
High dependence upon groundwater		
Muridae	<i>Hydromys chrysogaster</i>	water rat or rakali
Strong dependence upon groundwater		
Peramelidae	<i>Isodon obesulus fusciventer</i>	southern brown bandicoot or quenda
Macropodidae	<i>Macropus irma</i>	western brush wallaby
Muridae	<i>Rattus fuscipes</i>	southern bush rat
Low dependence upon groundwater		
Phalangeridae	<i>Trichosurus vulpecula</i>	brushtail possum
Burramyidae	<i>Cercartetus concinnus</i>	western pygmy possum
Tachyglossidae	<i>Tachyglossus aculeatus</i>	echidna
Dasyuridae	<i>Dasyurus geoffroii</i>	chuditch
Dasyuridae	<i>Sminthopsis dolichura</i>	little long-tailed dunnart
Dasyuridae	<i>Sminthopsis griseoventer</i>	grey-bellied dunnart
Tarsipedidae	<i>Tarsipes rostratus</i>	honey possum
Macropodidae	<i>Macropus fuliginosus</i>	western grey kangaroo

1035


1036

1037 Table 5: Susceptibility of extant terrestrial mammals to declining rainfall on the GGS.

Species	Relationship with Rainfall	Study
<i>Macropus fuliginosus</i>	Rainfall is the dominant influence on population dynamics in arid, semi-arid systems. Significant population declines with drought when females may cease breeding and mortality may exceed 40%. A short response lag (6 months) implies that survival, not breeding is impacted most.	Arnold and Steven 1988; Cairns <i>et al.</i> 2000; Caughley 1987; Caughley <i>et al.</i> 1984; Caughley <i>et al.</i> 1985; Coulson 2008
<i>Cercartetus concinnus</i>	Population sizes can fluctuate widely in dry regions. May be response to variability in rainfall and consequent food availability.	Carthew <i>et al.</i> 2008
<i>Tarsipes rostratus</i>	Low rainfall significantly influences abundance, and correlated with rainfall from 1-2 years previous. Reduction in numbers following low rainfall attributable to a decrease in nectar production.	Bradshaw <i>et al.</i> 2007; Bryant 2004; Wooller <i>et al.</i> 1998
<i>Isoodon obesulus fusciventer</i>	Substantial geographical variation in body size associated with habitat structure and associated with the amount of annual rainfall.	Cooper 1998; 2000a
<i>Rattus fuscipes</i>	Population fluctuations related to rainfall and the density of groundcover. High populations related to increased rainfall, drought associated with low numbers.	Lunney 2008; Lunney <i>et al.</i> 1987; Seebeck and Menkhorst 2000; Recher <i>et al.</i> 2009
<i>Hydromys chrysogaster</i>	<i>Hydromys chrysogaster</i> is found in fresh, brackish and saltwater wetlands. Indirect effect of rainfall based on water levels in wetlands and lakes.	Seebeck and Menkhorst 2000
<i>Tachyglossus aculeatus</i>	Indirect effect of rainfall on termites in arid areas. Rainfall, soil moisture affect termite availability-they move deeper if soil moisture is low.	Abensperg-Traun 1994

1038


1039



1040

1041 Figure 1. The remnant vegetation extent and DEC managed lands located within the GGS
 1042 study area. Fauna survey sites, including targeted mammal trapping areas are also
 1043 indicated.

1044



1045


1046 Figure 2. Long-term (1905-2007) annual rainfall for Wanneroo site 9105. (Source:

1047 Bureau of Meteorology.)

1048

1049

1050




1051

1052 Figure 3. Groundwater storage decline in the superficial aquifer since 1979. (Source:
1053 Department of Water.)

1054

1055



1056

1057

1058 Figure 4. The predicted impact of incremental declines in annual rainfall (based on the
 1059 long-term average annual rainfall of 807 mm / year) on the density of *Tarsipes rostratus*
 1060 capture rates (using the relationship between honey possum capture rates and rainfall
 1061 described in Wooller *et al.* (1998); capture rate = $0.199 \times \text{annual rainfall} + 2.8378$).
 1062 Annual rainfall records within the GGS (data from Bureau of Meteorology, Pearce RAAF
 1063 station number 009053) from 2006 – 2010 are indicated.